# AN OPERATOR INEQUALITY

Derming Wang

California State University–Long Beach

**1. Introduction.** In this paper inequalities for positive operators acting on a Hilbert space are considered. For positive operators $A$ and $B$, the following conjecture [1] was posed:

$$\text{If } 0 \le B \le A, \text{ then } (AB^2A)^{1/2} \le A^2. \tag{1}$$

This conjecture was answered affirmatively by Furuta [3]. Indeed, Furuta proved a more general inequality which contains inequality (1) as a special case:

$$\text{If } 0 \le B \le A, \text{ then } (A^r B^p A^r)^{1/q} \le A^{(p+2r)/q}$$
$$\text{for } p, r \ge 0, q \ge 1 \text{ with } p + 2r \le (1+2r)q.$$

Setting $p = q = 2$ and $r = 1$, Furuta's inequality becomes (1). More interestingly, if one sets $p = 2r$ and $q = 2$, Furuta's inequality becomes a generalization of (1):

$$\text{If } 0 \le B \le A, \text{ then } (A^r B^{2r} A^r)^{1/2} \le A^{2r} \text{ for } r \ge 0. \tag{2}$$

If $A$ and $B$ are positive invertible operators with $B \le A$, then it is known that $\log B \le \log A$. In [2], a result of Ando was rephrased: For positive invertible operators $A$ and $B$, $\log B \le \log A$ if and only if $(A^r B^{2r} A^r)^{1/2} \le A^{2r}$ holds for all $r \ge 0$. Thus, Ando's result also establishes (2) as a corollary.

In this paper we establish (2) directly by elementary means. Our approach is inspired by the work of Pedersen and Takesaki [5].

**2. Preliminary.** We are interested in (bounded, linear) operators acting on a Hilbert space $\mathbb{H}$ with inner product $< ., . >$. An operator $A$ is said to be self–adjoint if $A = A^*$, where $A^*$ is the adjoint of $A$. A self-adjoint operator $A$ is said to be positive, in notation $A \ge 0$ (or $0 \le A$), if $< Ax, x > \ge 0$ for every vector $x \in \mathbb{H}$. For positive operators $A$ and $B$

we write $A \geq B$ (or $B \leq A$) if $A - B \geq 0$. The relation "$\geq$" defines a partial order on the set of positive operators. The following properties of positive operators are well known:

(a) If $A \geq 0$, then $A^r \geq 0$ for every real number $r \geq 0$.

(b) If $A \geq 0$ and $A$ is invertible, then $A^{-1} \geq 0$.

(c) If $A, B \geq 0$, then $ABA \geq 0$. Thus, if $0 \leq B \leq A$, then $CBC \leq CAC$ for every $C \geq 0$.

(d) If $0 \leq B \leq A$, then $B^r \leq A^r$ for $0 \leq r \leq 1$. In particular, if $0 \leq B \leq A$, then $B^{1/2} \leq A^{1/2}$.

(e) In general, $0 \leq B \leq A$ does not necessarily imply $B^2 \leq A^2$.

(f) If $0 \leq T \leq I$, the identity operator, then $T^2 \leq I$.

Properties (c), (d) and (e) led the authors of [1] to pose the conjecture (1). Now we state a weakened version of a result of Pedersen and Takesaki [5] that is suitable for our purpose. For the sake of completeness, a proof is given.

Theorem A. Suppose $A$ and $B$ are positive operators. Also, assume $A$ is invertible. Then, if $(A^{1/2}BA^{1/2})^{1/2} \leq A$, then there is a unique positive operator $T$ satisfying $0 \leq T \leq I$ and $TAT = B$.

Proof. Let $T = A^{-1/2}(A^{1/2}BA^{1/2})A^{-1/2}$. Since $(A^{1/2}BA^{1/2})^{1/2} \leq A$, multiplying both sides of this inequality on the left and on the right by $A^{-1/2}$ yields $0 \leq T \leq I$. A simple calculation shows that $TAT = B$. This establishes the existence of $T$. To prove the uniqueness, assume $S$ is a positive operator with the property $SAS = B$. Since $SAS = TAT$, multiplying both sides of this equality on the left and on the right by $A^{1/2}$ we obtain $(A^{1/2}SA^{1/2})^2 = (A^{1/2}TA^{1/2})^2$. Now, taking square roots of both sides and then multiplying both sides on the left and on the right by $A^{-1/2}$ produces the desired $S = T$.

**3. The Main Result.** We are ready to present our proof of (2).

Lemma. If $0 \leq B \leq A$, then $(A^{2^{n-1}}B^{2^n}A^{2^{n-1}})^{1/2} \leq A^{2^n}$ for $n = 0, 1, 2, \ldots$.

Proof. First note that if the inequality can be proven with the extra assumption that the operator $A$ is invertible, then the result follows. For if $A$ is not invertible, then for any $\epsilon > 0$, the operator $A_\epsilon = A + \epsilon I$ is a positive invertible operator. If the inequality in the lemma is derived with $A_\epsilon$ in place of $A$, taking limits as $\epsilon$ tends to 0 will produce the desired inequality. Therefore, we may, without loss of generality, assume that $A$ is invertible.

Since $B \leq A$, $(A^{1/2}BA^{1/2})^{1/2} \leq A$. This proves the lemma for $n = 0$. Now, Theorem A implies there is a positive operator $T_1 \leq I$ such that $T_1AT_1 = B$. Thus,

$$AB^2A = A(T_1AT_1)^2A = (AT_1A)T_1^2(AT_1A) \leq (AT_1A)^2.$$

Taking square roots, we have $(AB^2A)^{1/2} \leq AT_1A \leq A^2$. This establishes the lemma for $n = 1$. Again, Theorem A implies there is a positive operator $T_2 \leq I$ such that $T_2A^2T_2 = B^2$. Similar arguments give $(A^2B^4A^2)^{1/2} \leq A^4$. This establishes the lemma for $n = 2$. It is now apparent that the lemma follows by induction.

Notice that for the case $n = 1$, the inequality of the Lemma is (1).

Theorem. If $0 \leq B \leq A$, then $(A^rB^{2r}A^r)^{1/2} \leq A^{2r}$ for $r \geq 0$.

Proof. For each $r \geq 0$, there is a smallest nonnegative integer $k$ such that $r/2^k \leq 1$. Let $A_1 = A^{r/2^k}$ and $B_1 = B^{r/2^k}$. We have $0 \leq B_1 \leq A_1$, $A_1^{2^k} = A^r$ and $B_1^{2^k} = B^r$. The result follows by applying the Lemma with $n = k + 1$ to the operators $A_1$ and $B_1$. This completes the proof.

We now use the Theorem to amplify Theorem A when $0 \leq B \leq A$.

Corollary. Suppose $0 \leq B \leq A$ with $A$ invertible. Then, for each $r \geq 0$ there is a unique operator $T_r$ satisfying $0 \leq T_r \leq I$ and $T_rA^rT_r = B^r$.

Proof. Let $T_r = A^{-r}(A^rB^{2r}A^r)^{1/2}A^{-r}$. Clearly we have $T_rA^rT_r = B^r$. Since $B \leq A$, the Theorem implies $0 \leq T_r \leq I$. Arguments similar to those of Theorem A show that the existence of $T_r$ is unique. The proof is complete.

In conclusion, we note that a result of Hansen [4] may be applied to show that the family $\{T_r\}_{r \geq 0}$ in the Corollary is a decreasing family.

## *References*


1. N. N. Chan and M. K. Kwong, "Hermitian Matrix Inequalities and a Conjecture," *American Mathematical Monthly*, 92 (1985), 533–541.
2. M. Fujii and E. Kamei, "Furuta's Inequality for Chaotic Order," *Mathematica Japonica*, 36 (1991), 603–606.
3. T. Furuta, "$A \geq B \geq 0$ Assures $(B^rA^pB^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0$, $p \geq 0$, $q \geq 1$ with $(1 + 2r)q \geq p + 2r$," *Proceedings of the American Mathematical Society*, 101 (1987), 85–88.
4. F. Hansen, "An Operator Inequality," *Mathematische Annalen*, 246 (1980), 249–250.
5. G. K. Pedersen and M. Takesaki, "The Operator Equation $THT = K$," *Proceedings of the American Mathematical Society*, 36 (1972), 311–312. æ